pregnated by the salt springs of the interior. The degree in which they require the presence of the mineral is various, some growing upon the beach, where the earth is saturated with salt, and others at the extreme edge of marshes, where the impregnation is much less powerful. With a few exceptions, they cannot long be cultivated in fresh earth, but soon decay when removed from their native marshes.

Maritime plants derive a peculiar character from their place of growth, which distinguishes them even when dry from other vegetables. The salt with which they are impregnated crystallizes on their surface in dry weather; and deliquesces so as to render them damp and supple, when the atmosphere is moist. These plants are troublesome in an herbarium from the facility with which they contract moisture from the atmosphere, and communicate it to the adjacent papers. The hay cut upon salt marshes often becomes extremely damp, and would be entirely spoiled, were it not for the antiseptic and preservative quality of the salt. The barilla of commerce is obtained by the combustion of maritime vegetables.

Many of these plants are thick and fleshy in their mode of growth, and differ remarkably in this respect from their co-species on dry ground. This is

particularly seen in Arenaria, Gerardia, Chenopodium, &c.

The vegetable which is the subject of this article is exclusively a maritime plant. It is one of the few ornamental species in our salt marshes, and is very conspicuous for its purple tops appearing among the grass in all the summer months. It varies from a few inches, to a foot and more in height.

This species has generally been considered a variety of the *Statice limonium*, which is a common plant in the salt marshes of Europe. Indeed, several of the maritime species of this genus approach each other so closely in their characters, that they have been considered the same by able botanists. The American plant, to which the name of *Caroliniana* was given by Walter in his Flora of Carolina, is distinguished from the European principally by its smaller flowers and plain or flat leaves. From the *Statice Gmelini*, an Asiatic species, it differs apparently still less in its general form.

The genus *Statice* belongs to the class *Pentandria* and order *Pentagynia*. Its natural orders are *Aggregatæ* of Linnæus and *Plumbagines* of Jussieu. It is characterized by *a calyx monophyllous, plaited* and *scarious. Petals five with*

JACOB BIGELOW *American Medical Botany* BOSTON, 1817-1820 CARY GRAPHIC ARTS COLLECTION, ROCHESTER INSTITUTE OF TECHNOLOGY Octavo